

ON THE GR-SPEEDS OF LIGHT AND PARTICLES (A REMEMBRANCE OF THINGS PAST)

ANGELO LOINGER AND TIZIANA MARSICO

ABSTRACT. There are no limits for the speeds of light and particles in general relativity (GR). Four examples illustrate this basic result, which is too often neglected.

PACS 04.20 – General relativity.

1. – The recent interesting experiments by the OPERA Collaboration concerning the neutrino velocity [1] have given rise indirectly to renewed discussions about the velocities of light and particles in general relativity. – In this Note we emphasize the following fundamental concepts: *i*) In GR the light velocity can have *any* value between zero and infinite, according to the values of the metric tensor; *ii*) *Idem* for the particle velocity, which however must always be lower than the light velocity in the *same* pseudo-Riemannian manifold, but it can be greater than c (the light velocity *in vacuo*, *i.e.* in the absence of any field) in some instances.

We shall illustrate these statements with four simple examples.

2. – A preliminary consideration. In an axiomatic treatment of GR – see, *e.g.*, Hilbert [2] – the theory is divided in two distinct parts. The first part is devoted to the properties of the Einsteinian field equations, without any indication of the Riemannian or pseudo-Riemannian character of the interested manifolds.

In the second part, according to the *Erfahrung*, Hilbert specifies the nature of the interval ds^2 , which can be time-like, light-like, or space-like. The spacetime manifolds of physical interest must have a well-defined pseudo-Riemannian character.

The points *i*) and *ii*) of sect.1 are implicitly contained in this axiom, *i.e.* they are a logical consequence of it.

3. – Let us consider the Einsteinian stationary field which exists in a reference frame uniformly rotating [3], [4]. The interval ds^2 is:

$$(1) \quad ds^2 = (c^2 - \Omega^2 r^2) dt^2 - 2\Omega r^2 d\varphi dt - d\sigma^2 \quad , \quad (c^2 > \Omega^2 r^2) \quad ,$$

$$(1') \quad d\sigma^2 \equiv dr^2 + r^2 d\varphi^2 + dz^2 \quad ,$$

where Ω is the angular speed of rotation, and r, φ, z are cylindrical space-coordinates. Light motions are characterized by $ds^2 = 0$. If Ω is *small*, we can keep only the terms of *first* order in $\Omega r^2/c$. We get:

$$(2) \quad dt = \frac{d\sigma}{c} \pm \frac{2\Omega}{c^2} \left(\frac{1}{2} r^2 d\varphi \right) \quad ,$$

where the double sign (\pm) corresponds to the two directions of the light-ray motion.

If we put $r = a_1$, the Earth's equatorial radius ($a_1 = 6,378,137$ m), and $\Omega = \omega_E$, the terrestrial rotation speed ($\omega_E = 7.2921151467 \times 10^{-5}$ rad/s), for a full turn (within an optical fibre) of a light-ray around the equator we have:

$$(3) \quad \Delta t = \frac{2\pi a_1}{c} \pm \frac{2\omega_E}{c^2} \pi a_1^2 \quad ;$$

since $\pi a_1^2 = 1.27802 \times 10^{14} \text{ m}^2$, and $2\omega_E/c^2 = 1.6227 \times 10^{-21} \text{ m}^{-2}\text{s}$, the second term at the right-side of (3) – which gives a so-called Sagnac effect – is equal to 207.4 ns.

The light velocity w is:

$$(4) \quad w = \frac{2\pi a_1}{\Delta t} = c \mp \omega_E a_1 \quad ; \quad (\omega_E a_1 = 0.4651011 \text{ km/s}) \quad ;$$

we see that if the motions of the light-ray and of the reference frame happen in opposite directions, the light velocity is *greater* than c .

It is useful to compute w as an instantaneous speed. From $ds^2 = 0$, we obtain:

$$(5) \quad \frac{dt}{d\varphi} = \pm \frac{\omega_E a_1^2}{c^2} + \left(\frac{\omega_E^2 a_1^4}{c^4} + \frac{a_1^2}{c^2} \right)^{1/2} \approx \pm \frac{\omega_E a_1^2}{c^2} + \frac{a_1}{c} \quad ;$$

from which:

$$(6) \quad a_1 \frac{d\varphi}{dt} \approx \frac{c^2}{c \pm \omega_E a_1} \quad ;$$

eq. (6) coincides with eq. (4) if we neglect $\omega_E^2 a_1^2$.

4. – With a good approximation, the Einsteinian metric near the Earth can be written as follows (see, *e.g.*, [4]):

$$(7) \quad ds^2 \approx \left(1 + \frac{2V}{c^2} \right) c^2 dt^2 - 2\omega_E r^2 \sin^2 \vartheta d\varphi dt - \left(1 - \frac{2V}{c^2} \right) d\sigma^2 \quad ,$$

$$(7') \quad d\sigma^2 \equiv dr^2 + r^2 (d\vartheta^2 + \sin^2 \vartheta d\varphi^2) \quad ,$$

if we use isotropic polar coordinates r, ϑ, φ .

$$(8) \quad V \approx -\frac{GM_E}{r} \left[1 - \mathcal{J}_2 \left(\frac{a_1}{r} \right)^2 P_2(\cos \vartheta) \right] \quad ,$$

where: M_E is the Earth's mass, \mathcal{J}_2 is Earth's quadrupole moment coefficient, P_2 is the Legendre polynomial of degree 2; ($GM_E = 3.986004418 \times 10^{14} \text{ m}^3 \text{ s}^{-2}$; $\mathcal{J}_2 = 1.0826300 \times 10^{-3}$). From $ds^2 = 0$, we have, keeping only the terms of *first* order in $\omega_E r^2/c$ and V/c^2 :

$$(9) \quad c dt \approx \frac{d\sigma \pm \omega_E r^2 \sin^2 \vartheta d\varphi/c}{1 + 2V/c^2} \quad ;$$

for a full turn of a light-ray around the equator, we get [$P_2(\cos \pi/2) = -1/2$]:

$$(10) \quad a_1 \frac{d\varphi}{dt} \approx \frac{c^2}{c \pm \omega_E a_1} \left[1 - \frac{2|V(a_1)|}{c^2} \right] \quad ,$$

where

$$(11) \quad |V(a_1)| \approx \frac{GM_E}{a_1} (1 + 0.54 \times 10^{-3}) \approx \frac{GM_E}{a_1} \quad ,$$

from which:

$$(12) \quad \begin{aligned} a_1 \frac{d\varphi}{dt} &\approx \frac{c^2}{c \pm \omega_E a_1} \left(1 - \frac{2GM_E/c^2}{a_1} \right) \approx \\ &\approx (c \mp \omega_E a_1) \times (1 - 1.4 \times 10^{-9}) \approx \\ &\approx (c \mp 0.4651011) \text{ km/s} \quad . \end{aligned}$$

A result which is essentially coincident with that of the mere rotation.

5. – In Newton's theory, Laplace equation $\nabla^2 U = 0$ admits the two solutions $U = \pm GM/r$ for a material point. Analogous situation in GR. Let us start from the standard (Hilbert-Droste-Weyl) form of solution to Schwarzschild's problem of the gravitational field created by a point-mass M :

$$(13) \quad ds^2 = \left(1 - \frac{2m}{r} \right) c^2 dt^2 - \left(1 - \frac{2m}{r} \right)^{-1} dr^2 - r^2 (d\vartheta^2 + \sin^2 \vartheta d\varphi^2) \quad ,$$

where $m \equiv GM/c^2$ is an integration constant, that is specified in this way because we wish to have the Newton solution at great values of r . Now, the Einstein equations $R_{jk} = 0$, ($j, k = 0, 1, 2, 3$), admit also the following solution, which has a mere mathematical meaning (even if $ds^2 \geq 0$):

$$(14) \quad ds^2 = \left(1 + \frac{k}{r} \right) c^2 dt^2 - \left(1 + \frac{k}{r} \right)^{-1} dr^2 - r^2 (d\vartheta^2 + \sin^2 \vartheta d\varphi^2) \quad ,$$

where $k > 0$ is an integration constant; the condition $ds^2 = 0$ implies:

$$(15) \quad \left| \frac{dr}{dt} \right|_{\text{LIGHT}} = c \left(1 + \frac{k}{r} \right) > c \quad ;$$

and for the radial geodesic of a test-particle we have:

$$(16) \quad \left| \frac{dr}{dt} \right|_{\text{PARTICLE}} = c \left(1 + \frac{k}{r} \right) \left[1 - |A| \left(1 + \frac{k}{r} \right) \right] \quad ,$$

where A is an integration constant (for $A = 0$, we re-obtain eq. (15)). If $|A|$ is sufficiently small, *the particle velocity is greater than c for all the values of r greater than a suitable value.*

6. – The metric of Friedmann’s model of universe corresponding to a positive space curvature is represented by the following interval ds^2 (where x, y, z, r are dimensionless coordinates which give labels to the galaxies):

$$(17) \quad ds^2 = c^2 dt^2 - A^2(r^2) F^2(t) d\sigma^2 \quad ; \quad (r^2 = x^2 + y^2 + z^2) \quad ,$$

$$(17') \quad d\sigma^2 = dx^2 + dy^2 + dz^2 \quad ,$$

$$(17'') \quad A(r^2) = 1 + (r^2/4) \quad .$$

The function $F(t)$ – *scale factor* – is determined by Einstein field equations (with a “dust-like” matter tensor), and is a usual cycloid. Thus, we have a cosmic periodic oscillation between $F = 0$ and $F = F_{\text{max}}$.

From $ds^2 = 0$, we get:

$$(18) \quad \left| \frac{d\sigma}{dt} \right|_{\text{LIGHT}} = \frac{c}{A(r^2) F(t)} \quad ;$$

we see that the light velocity is infinite at every big-bang and at every big-crunch. –

Acknowledgment. We are very grateful to Dr. L. Stanco of the OPERA Collaboration for the epistolary discussions about several passages of the papers quoted in [1].

REFERENCES

- [1] T. Adam *et alii*, *arXiv:1109.4897 v2* [hep-ex] Nov 18, 2011. See also: E. Kiritsis and F. Nitti, <http://operaweb.lngs.infn.it:2080/Opera/publicnotes/note136.pdf> – G. Brunetti, *PhD thesis*, in joint supervision from Université Claude Bernard Lyon-1 and Università di Bologna, 2011 – http://operaweb.lngs.infn.it:2080/Opera/ptb/theses/Brunetti-Giulia_phdthesis.pdf
- [2] D. Hilbert, *Mathem. Annalen*, **92** (1924) 1.
- [3] L. Landau et E. Lifchitz, *Théorie du Champ* (Éditions Mir, Moscou) 1966, sects. **84** and **90**.
- [4] N. Ashby, *Living Reviews in Relativity* (2003-1) <http://www.livingreviews.org>.

A.L. – DIPARTIMENTO DI FISICA, UNIVERSITÀ DI MILANO, VIA CELORIA, 16 - 20133 MILANO (ITALY)

T.M. – LICEO CLASSICO “G. BERCHET”, VIA DELLA COMMENDA, 26 - 20122 MILANO (ITALY)

E-mail address: angelo.loinger@mi.infn.it

E-mail address: martiz64@libero.it